

On the monodromy group of confluent linear equations

Alexey GLUTSYUK

February 1, 2008

Dedicated to Yu.S.Ilyashenko on the occasion of his 60th birthday.

Abstract

We consider a linear analytic ordinary differential equation with complex time having a nonresonant irregular singular point. We study it as a limit of a generic family of equations with confluent Fuchsian singularities.

In 1984 V.I. Arnold asked the following question: is it true that some operators from the monodromy group of the perturbed (Fuchsian) equation tend to Stokes operators of the nonperturbed irregular equation? Another version of this question was also independently proposed by J.-P. Ramis in 1988.

We consider the case of Poincaré rank 1 only. We show (in dimension two) that generically no monodromy operator tends to a Stokes operator; on the other hand, in any dimension commutators of appropriate noninteger powers of the monodromy operators around singular points tend to Stokes operators.

2000 Math. Subj. Class. 34M35 (34M40)

Key words and phrases: Linear equation, irregular singularity, Stokes operators, Fuchsian singularity, monodromy, confluence.

Table of contents.

1. Introduction.
 - 1.1. Brief statements of results, the plan of the paper and the history.
 - 1.2. Analytic classification of irregular equations. Canonical solutions and Stokes operators.
 - 1.3. Historical overview.
2. Main results. Stokes operators and limit monodromy.
 - 2.1. Stokes operators as limit transition operators between monodromy eigenbases.
 - 2.2. Stokes operators as limits of commutators of appropriate powers of the monodromy operators.
 - 2.3. The case of higher Poincaré rank.
3. Convergence of the commutators to Stokes operators. Proof of Theorem 2.16.
 - 3.1. Properties of the monodromy and the transition operators. The plan of the proof of Theorem 2.16.

- 3.2. The upper-triangular element of the transition matrix. Proof of (3.3).
- 3.3. Commutators of operators with asymptotically common eigenline. Proof of Lemma 3.5.
- 3.4. Convergence of the commutators to Stokes operators: the higher-dimensional case.
- 4. Generic divergence of monodromy operators along degenerating loops.
 - 4.1 The statement of the divergence Theorem.
 - 4.2. Projectivization. The scheme of the proof of Theorem 4.6.
 - 4.3. Divergence of word of projectivizations. Proof of Lemma 4.10.

References.

1 Introduction

1.1 Brief statements of results, the plan of the paper and the history

Consider a linear analytic ordinary differential equation

$$\dot{z} = \frac{A(t)}{t^{k+1}}z, \quad z \in \mathbb{C}^n, \quad |t| \leq 1, \quad k \in \mathbb{N} \quad (1.1)$$

with a nonresonant irregular singularity of order (the Poincaré rank) k at 0 (or briefly, an irregular equation). This means that $A(t)$ is a holomorphic matrix function such that the matrix $A(0)$ has distinct eigenvalues (denote them by λ_i). Then the matrix $A(0)$ is diagonalizable, and without loss of generality we suppose that it is diagonal.

1.1 Definition Two equations of type (1.1) are analytically (formally) equivalent, if there exists a change $z = H(t)w$ of the variable z , where $H(t)$ is a holomorphic invertible matrix function (respectively, a formal invertible matrix power series), that transforms one equation into the other.

The analytic classification of irregular equations (1.1) is well-known ([2], [3], [9], [10], [16]): the complete system of invariants for analytic classification consists of a formal normal form (1.4) and Stokes operators (1.6) defined in 1.2; the latter are linear operators acting in the solution space of (1.1) comparing appropriate "sectorial canonical solution bases".

On the other hand, an irregular equation (1.1) can be regarded as a result of *confluence of Fuchsian singular points* (recall that a Fuchsian singular point of a linear equation is a first order pole of its right-hand side). Namely, consider a deformation

$$\dot{z} = \frac{A(t, \varepsilon)}{f(t, \varepsilon)}z, \quad f(t, \varepsilon) = \prod_{i=0}^k (t - \alpha_i(\varepsilon)), \quad (1.2)$$

of equation (1.1) that splits the irregular singular point 0 of the nonperturbed equation into $k+1$ Fuchsian singularities $\alpha_i(\varepsilon)$ of the perturbed equation, i.e., $\alpha_i(\varepsilon) \neq \alpha_j(\varepsilon)$ for $i \neq j$. The family (1.2) depends on a parameter $\varepsilon \in \mathbb{R}_+ \cup 0$, $f(t, 0) \equiv t^{k+1}$, $A(t, 0) \equiv A(t)$.

The *monodromy group* of a Fuchsian equation acts linearly in its solution space by analytic extensions of solutions along closed loops. The analytic equivalence class of a generic Fuchsian

equation is completely determined by the local types of its singularities and the action of its monodromy group.

Everywhere below by M_i we denote the monodromy operator of the perturbed equation (1.2) along a loop going around the singular point α_i (the choice of the corresponding loops will be specified later). The monodromy group of the perturbed equation is generated by appropriately chosen operators M_i .

In 1984 V.I. Arnold proposed the following question. Consider a generic deformation (1.2). Is there an operator

$$M_{i_l}^{d_l} \dots M_{i_1}^{d_1} \quad (1.3)$$

from the monodromy group of the perturbed equation that converges to a Stokes operator of the nonperturbed equation?

A version of this question was proposed independently by J.-P. Ramis in 1988.

It appears that already in the simplest case of dimension 2 and Poincaré rank $k = 1$ generically *each operator from the monodromy group (except for that along a circuit (and its powers) around both singularities) tends to infinity* (Theorem 4.6 in Section 4), so, no one tends to a Stokes operator.

In other terms, generically, no word (1.3) with $d_i \in \mathbb{Z}$ tends to a Stokes operator. But if $k = 1$, then appropriate words (1.3) with *noninteger* exponents d_i tend to Stokes operators (Theorem 2.16 in 2.2).

The previous question and its nonlinear analogues were studied by J.-P. Ramis, B. Khesin, A. Duval, C. Zhang, J. Martinet, the author and others (see the historical overview in 1.3). It was proved by the author in [6] that appropriate branches of the eigenfunctions of the monodromy operators M_i of the perturbed equation tend to appropriate canonical solutions of the nonperturbed equation (Theorem 2.5 in 2.1). In the case of Poincaré rank $k = 1$ this implies (Corollary 2.6 in 2.1) that Stokes operators of the nonperturbed equation are limits of transition operators between appropriate eigenbases of the monodromy operators M_i . This Corollary has a generalization for higher Poincaré rank [6].

The proofs of the results of the present paper are based on the previously mentioned results from [6], which are recalled in 2.1.

In 1.2 we recall the analytic classification of irregular equations (1.1) and the definitions of sectorial canonical solution bases and Stokes operators.

In 2.2 we state Theorem 2.16 on convergence of appropriate word (1.3) with noninteger exponents d_i to a Stokes operator in the case of Poincaré rank $k = 1$. Its proof is given in Section 3. The corresponding exponents d_i do not depend on the choice of deformation. In fact, in the case of the higher Poincaré rank $k = 2$ and $n = 2$ one can also prove a similar statement, but now the choice of the corresponding exponents d_i will depend on the choice of deformation. The latter case will be discussed in 2.3.

In Section 4 in the case, when $k = 1$, $n = 2$, for a typical nonperturbed equation (1.1) we prove the divergence of the operators from the monodromy group of the perturbed equation (except for the monodromy along a circuit around both singularities and its powers).

1.2 Analytic classification of irregular equations. Canonical solutions and Stokes operators

Let (1.1) be an irregular equation, λ_i , $i = 1, \dots, n$, be the eigenvalues of the corresponding matrix $A(0)$.

One can ask the question: is it true that the variables $z = (z_1, \dots, z_n)$ in the equation can be separated, more precisely, that (1.1) is analytically equivalent to a direct sum of one-dimensional linear equations, i.e., a linear equation with a diagonal matrix function in the right-hand side? Generically, the answer is “no”. At the same time any irregular equation (1.1) is formally equivalent to a unique direct sum of the type

$$\begin{cases} \dot{w}_i = \frac{b_i(t)}{t^{k+1}} w_i, \\ i = 1, \dots, n, \end{cases} \quad (1.4)$$

where $b_i(t)$ are polynomials of degrees at most k , $b_i(0) = \lambda_i$. The normalizing series bringing (1.1) to (1.4) is unique up to left multiplication by constant diagonal matrix. The system (1.4) is called the *formal normal form* of (1.1) ([2], [3], [9], [10], [16]).

Generically the normalizing series diverges. At the same time there exists a finite covering $\bigcup_{j=0}^N S_j$ of a punctured neighborhood of zero in the t -line by radial sectors S_j (i.e., those with the vertex at 0) that have the following property. There exists a unique change of variables $z = H_j(t)w$ over each S_j that transforms (1.1) to (1.4), where $H_j(t)$ is an analytic invertible matrix function on S_j that can be C^∞ -smoothly extended to the closure $\overline{S}_j$ of the sector so that its asymptotic Taylor series at 0 coincides with the normalizing series. The previous statement on existence and uniqueness of sectorial normalization holds in any good sector (see the two following Definitions); the covering consists of good sectors ([2], [3], [9], [10], [16]).

Case $k = 1$, $n = 2$, $\lambda_1 - \lambda_2 \in \mathbb{R}$.

1.2 Definition A sector in $\mathbb{C}$ with the vertex at 0 is said to be *good*, if it contains only one imaginary semiaxis $i\mathbb{R}_\pm$, and its closure does not contain the other one (see Fig.1).

General case.

1.3 Definition (see, e.g., [9]). Let $k \in \mathbb{N}$, $\Lambda = \{\lambda_1, \dots, \lambda_n\} \subset \mathbb{C}$ be a n -ple of distinct numbers, t be the coordinate on $\mathbb{C}$. For a given pair $\lambda_i \neq \lambda_j$ the rays in $\mathbb{C}$ starting at 0 and forming the set $\text{Re}(\frac{\lambda_j - \lambda_i}{t^k}) = 0$ are called the (k, Λ) - *imaginary dividing rays* corresponding to the pair (λ_i, λ_j) . A radial sector is said to be (k, Λ) - *good*, if for any pair (λ_i, λ_j) , $j \neq i$, it contains exactly one corresponding imaginary dividing ray and so does its closure.

1.4 Remark In the case, when $k = 1$, $n = 2$, $\lambda_1 - \lambda_2 \in \mathbb{R}$, the imaginary dividing rays are the imaginary semiaxes, and the notions of good sector and (k, Λ) - good sector coincide.

1.5 Remark The ratio $\frac{w_i}{w_j}(t)$ of solutions of equations from (1.4) tends to either zero or infinity, as t tends to zero along a ray distinct from the imaginary dividing rays corresponding to the pair (λ_i, λ_j) . Its limit changes exactly when the ray under consideration jumps over one of the latter imaginary dividing rays.

We consider a covering $\bigcup_{j=0}^N S_j$ of a punctured neighborhood of zero by good (or (k, Λ) -good) sectors numerated counterclockwise and put $S_{N+1} = S_0$. The standard splitting of the normal form (1.4) into the direct sum of one-dimensional equations defines a canonical base

in its solutions space (uniquely up to multiplication of the base functions by constants) with a diagonal fundamental matrix. Denote the latter fundamental matrix by

$$W(t) = \text{diag}(w_1, \dots, w_n).$$

Together with the normalizing changes H_j in S_j , it defines the canonical bases $(f_{j1}, \dots, f_{jn})$ in the solution space of (1.1) in the sectors S_j with the fundamental matrices

$$Z^j(t) = H_j(t)W(t), \quad j = 0, \dots, N+1, \quad (1.5)$$

where for any $j = 0, \dots, N$ the branch ("with the index $j+1$ ") of the fundamental matrix $W(t)$ in S_{j+1} is obtained from that in S_j by the counterclockwise analytic extension for any $j = 0, \dots, N$. (We put $S_{N+1} = S_0$. The corresponding branch of W "with the index $N+1$ " is obtained from that "with the index 0" by the right multiplication by the monodromy matrix of the formal normal form (1.4).) In a connected component of the intersection $S_j \cap S_{j+1}$ there are two canonical solution bases coming from S_j and S_{j+1} . Generically, they do not coincide. The transition between them is defined by a constant matrix C_j :

$$Z^{j+1}(t) = Z^j(t)C_j. \quad (1.6)$$

The transition operators (matrices C_j) are called *Stokes operators (matrices)* (see [2], [3], [9], [10], [16]). The nontriviality of Stokes operators yields the obstruction to analytic equivalence of (1.1) and its formal normal form (1.4).

1.6 Remark Stokes matrices (1.6) are well-defined up to simultaneous conjugation by one and the same diagonal matrix.

1.7 Example Let $k = 1, n = 2$. In this case without loss of generality we assume that $\lambda_1 - \lambda_2 \in \mathbb{R}_+$ (one can achieve this by linear change of the time variable). Then the above covering consists of two sectors S_0 and S_1 (Figure 1). The former contains the positive imaginary semiaxis and its closure does not contain the negative one; the latter has the same properties with respect to the negative (respectively, positive) imaginary semiaxis. There are two components of the intersection $S_0 \cap S_1$. So, in this case we have a pair of Stokes operators. *The Stokes matrices (1.6) are unipotent*: the one corresponding to the left intersection component is lower-triangular; the other one is upper-triangular ([2], [3], [9], [10], [16]).

1.8 Remark Stokes operators of an irregular equation (1.1) with a diagonal matrix in the right-hand side are identity operators. In this case (1.1) is analytically equivalent to its formal normal form. In general, *two irregular equations are analytically equivalent, if and only if they have the same formal normal form and the corresponding Stokes matrix tuples are obtained from each other by simultaneous conjugation by one and the same diagonal matrix*, cf. the previous Remark. Thus, formal normal form and Stokes matrix tuple taken up to the previous conjugation present the complete system of invariants for analytic classification of irregular equations (see [2], [3], [9], [10], [16]).




Figure 1

1.3 Historical overview

Earlier in 1919 R. Garnier [5] had studied some particular deformations of some class of linear equations with nonresonant irregular singularity. He obtained some analytic classification invariants for these equations by studying their deformations. The complete system of analytic classification invariants (Stokes operators and formal normal form) for general irregular differential equations was obtained later in 1970-th years in the papers by Jurkat, Lutz, Peyerimhoff [10], Sibuya [16] and Balser, Jurkat, Lutz [3]. Later Jurkat, Lutz and Peyerimhoff had extended their results to some resonant cases [11]. It is well-known that the monodromy operators of a linear ordinary differential equation belong to its Galois group (see [9], [14]). In 1985 J.-P. Ramis have proved that the Stokes operators also belong to the Galois group ([14], see also [9]). In 1989 he considered the classical confluent family of hypergeometric equations and proved convergence of appropriate branches of monodromy eigenfunctions of the perturbed equation to canonical solutions of the nonperturbed one by direct calculation [15]. In the late 1980-th years B. Khesin also proved a version of this statement, but his result was not published. In 1991 A. Duval [4] proved this statement for the biconfluent family of hypergeometric equations (where the nonperturbed equation is equivalent to Bessel equation) by direct calculation. In 1994 C. Zhang [17] had obtained the expression of Garnier's invariants via Stokes operators (for the class of irregular equations considered by Garnier).

The conjecture saying that Stokes operators are limit transition operators between monodromy eigenbases of the perturbed equation was firstly proposed by A.A. Bolibrukh in 1996. It was proved by the author in [6]. Later this result was extended to a generic resonant case [8].

Nonlinear analogues of the previous statements for parabolic mappings (i.e., one-dimensional conformal mappings tangent to identity) and their Écalle-Voronin moduli, saddle-node singularities of two-dimensional holomorphic vector fields and their Martinet-Ramis invariants (sectorial central manifolds in higher dimensions) were obtained by the author in [7]. Generalizations and other versions of the statement on parabolic mappings were recently

obtained in the joint paper [12] by P.Mardesic, R.Roussarie, C.Rousseau, and in two unpublished joint papers by the following authors: 1) X.Buff and Tan Lei; 2) A.Douady, Francisco Estrada, P.Sentenac.

A particular case of the result from [7] concerning parabolic mappings (analogous to the previously mentioned statements on linear equations) was obtained by J.Martinet [13].

2 Main results. Stokes operators and limit monodromy

Everywhere below (whenever the contrary is not specified) we consider that the (nonperturbed) irregular equation under consideration has Poincaré rank $k = 1$. In the present Section we recall the statements from [6] expressing the Stokes operators as limit transition operators between monodromy eigenbases of the confluent Fuchsian equation (Theorems 2.5, 2.11 and Corollary 2.6 in Subsection 2.1). In 2.2 we state the results expressing the Stokes operators as limits of some words (1.3) of noninteger powers of monodromy operators (Theorem 2.16). In 2.3 we discuss the extension of these results to the case of higher Poincaré rank in dimension two.

2.1 Stokes operators as limit transition operators between monodromy eigenbases

We formulate the result from the title of the Subsection firstly in the case, when $k = 1$, $n = 2$, and then in the general case.

Case $n = 2$, $k = 1$. Let λ_i , $i = 1, 2$, be the eigenvalues of the matrix $A(0)$. Without loss of generality we assume that $\lambda_1 - \lambda_2 \in \mathbb{R}_+$: one can achieve this by linear change of the time variable.

We consider a deformation of (1.1),

$$\dot{z} = \frac{A(t, \varepsilon)}{f(t, \varepsilon)} z, \quad f(t, \varepsilon) = (t - \alpha_0(\varepsilon))(t - \alpha_1(\varepsilon)), \quad f(t, 0) \equiv t^2, \quad A(t, 0) = A(t), \quad (2.1)$$

where $A(t, \varepsilon)$ and $f(t, \varepsilon)$ depend continuously on a parameter $\varepsilon \geq 0$ so that $\alpha_0(\varepsilon) \neq \alpha_1(\varepsilon)$ for $\varepsilon > 0$. Without loss of generality we assume that $\alpha_0 + \alpha_1 \equiv 0$. We formulate the statement from the title of the Subsection for a generic deformation (2.1), see the following Definition.

2.1 Definition A family of quadratic polynomials $f(t, \varepsilon)$ depending continuously on a non-negative parameter ε , $f(t, 0) \equiv t^2$, with roots $\alpha_i(\varepsilon)$, $i = 0, 1$, $\alpha_0 + \alpha_1 \equiv 0$, is said to be *generic*, if $\alpha_0(\varepsilon) \neq \alpha_1(\varepsilon)$ for $\varepsilon \neq 0$, and the line passing through $\alpha_0(\varepsilon)$ and $\alpha_1(\varepsilon)$ intersects the real axis by angle bounded away from 0 uniformly in ε . A family (2.1) of *linear equations* with $n = 2$, $k = 1$, $\lambda_1 - \lambda_2 \in \mathbb{R}_+$ is said to be *generic*, if so is the corresponding family of polynomials $f(t, \varepsilon)$.

2.2 Definition (see, e.g., [2]). A singular point t_0 of a linear analytic ordinary differential equation $\dot{z} = \frac{B(t)}{t-t_0} z$ is said to be *Fuchsian*, if it is a first order pole of the right-hand side (i.e., the corresponding matrix function $B(t)$ is holomorphic at t_0). The *characteristic numbers* of a Fuchsian singularity are the eigenvalues of the corresponding residue matrix $B(t_0)$ (which are equal to the logarithms divided by $2\pi i$ of the eigenvalues of the corresponding monodromy operator).

2.3 Remark A family (2.1) of linear equations is generic, if and only if the difference of the characteristic numbers at $\alpha_0(\varepsilon)$ (or equivalently, at $\alpha_1(\varepsilon)$) of the perturbed equation is not real for small ε and moreover has argument bounded away from $\pi\mathbb{Z}$ uniformly in ε small enough. The latter condition implies that the monodromy operator of the perturbed equation around each singular point α_i has distinct eigenvalues (moreover, their modules are distinct), and hence, a well-defined eigenbase in the solution space (for small ε).

The singularities of the perturbed equation from a generic family have imaginary parts of constant (and opposite) signs (by definition). Without loss of generality everywhere below we consider that

$$\operatorname{Im} \alpha_0 > 0, \operatorname{Im} \alpha_1 < 0, \text{ see Fig. 2.}$$




Figure 2

2.4 Definition Let (2.1) be a generic family of linear equations (see the previous Definition) whose singularity families satisfy the previous inequalities. Let S_j , $j = 0, 1$, be a pair of good sectors in the t -line (see Definition 1.2) such that for any ε small enough $\alpha_j(\varepsilon) \in S_j$, $j = 0, 1$, $i\mathbb{R}_+ \subset S_0$, $i\mathbb{R}_- \subset S_1$ (see Fig. 1). The sector S_j is said to be *the sector associated to the singularity family α_j* , $j = 0, 1$.

We show that appropriate branches of the eigenfunctions of the monodromy operator M_i around α_i of the perturbed equation converge to canonical solutions of the nonperturbed equation in the corresponding sector S_i . This will imply the statement from the title of the Subsection.

To formulate the latter statement precisely, consider the auxiliary domain

$$S'_i = S_i \setminus [\alpha_0(\varepsilon), \alpha_1(\varepsilon)], \quad (2.2)$$

which is simply-connected, and the canonical branches of the monodromy eigenfunctions on the domain S'_i . In more details, consider a small circle going around α_i and take a base point on it outside the segment $[\alpha_0(\varepsilon), \alpha_1(\varepsilon)]$. In the space of local solutions of the perturbed equation at the base point consider the monodromy operator M_i acting by the analytic extension of a solution along the circle from the base point to itself in the counterclockwise direction. The eigenfunctions of M_i have well-defined branches (up to multiplication by

constants) in the corresponding disc with the segment $[\alpha_0(\varepsilon), \alpha_1(\varepsilon)]$ deleted. Their immediate analytic extension yields their canonical branches on S'_i . In other terms, we identify the space of local solutions with the space of solutions on S'_i by immediate analytic extension, consider M_i as an operator acting in the latter space and take its eigenfunctions.

The canonical basic solutions of the nonperturbed equation are numerated by the indices 1 and 2, which correspond to the eigenvalues λ_1, λ_2 of $A(0)$. To state the results previously mentioned, let us define the analogous numeration of the monodromy eigenfunctions at $\alpha_i(\varepsilon)$. The monodromy eigenfunctions are numerated by the characteristic numbers (see Definition 2.2) of the corresponding singularity. The latter are proportional to the eigenvalues of the matrix $A(\alpha_i(\varepsilon), \varepsilon)$, which tend to λ_1 and λ_2 , as $\varepsilon \rightarrow 0$. This induces the numeration of the monodromy eigenfunctions by the indices 1 and 2 corresponding to the limit eigenvalues λ_1 and λ_2 .

2.5 Theorem (see [6]). *Let (2.1) be a generic family of linear ordinary differential equations (see Definition 2.1), $\alpha_i(\varepsilon)$ be its singularity family, S_i be the corresponding sector (see the previous Definition), S'_i be the domain (2.2). Consider the eigenbase on S'_i of the monodromy operator of the perturbed equation around $\alpha_i(\varepsilon)$. The appropriately normalized eigenbase (by multiplication of the basic functions by constants) converges to the canonical solution base (1.5) on S_i of the nonperturbed equation.*

2.6 Corollary (see [6]). *Let (2.1) be a generic linear equation family (see Definition 2.1), α_i be its singularity families, S_i be the corresponding sectors (see the previous Definition) chosen to cover a punctured neighborhood of zero, S'_i be the corresponding domains (2.2). Let C_0, C_1 be the corresponding Stokes matrices (1.6) of the nonperturbed equation in the left (respectively, right) component of the intersection $S_0 \cap S_1$. Consider the eigenbase on S'_i of the monodromy operator of the perturbed equation around $\alpha_i(\varepsilon)$. Denote by $Z_\varepsilon^i(t)$ the fundamental matrix of this eigenbase. Let $C_0(\varepsilon)$ ($C_1(\varepsilon)$) be the transition matrix between the monodromy eigenbases $Z_\varepsilon^i(t)$, $i = 0, 1$, in the left (respectively, right) component of the intersection $S'_0 \cap S'_1$:*

$$Z_\varepsilon^1(t) = Z_\varepsilon^0(t)C_0(\varepsilon) \text{ for } \operatorname{Re} t < 0; \quad Z_\varepsilon^0(t) = Z_\varepsilon^1(t)C_1(\varepsilon) \text{ for } \operatorname{Re} t > 0. \quad (2.3)$$

For any $j = 0, 1$ and appropriately normalized monodromy eigenbases Z_ε^i , $i = 0, 1$ (the normalization of Z_ε^0 (only) depends on the choice of j) $C_j(\varepsilon) \rightarrow C_j$, as $\varepsilon \rightarrow 0$.

Case $k = 1$, n is arbitrary. To state the analogues of Theorem 2.5 and Corollary 2.6 in this more general case, let us firstly extend the notions of a generic family of linear equations and a sector associated to a singularity family.

2.7 Definition Let $n, k \in \mathbb{N}$, $n \geq 2$, $\Lambda = (\lambda_1, \dots, \lambda_n)$ be a set of n distinct complex numbers, $\lambda_i \neq \lambda_j$ be a pair of them. A ray in $\mathbb{C}$ starting at 0 is called a (k, Λ) - *real dividing ray* associated to the pair (λ_i, λ_j) , if for any t lying in this ray $\operatorname{Im} \frac{\lambda_i - \lambda_j}{t^k} = 0$. (Or equivalently, it is a ray bisecting an angle between two neighbor imaginary dividing rays (see Definition 1.3) associated to (λ_i, λ_j) .)

2.8 Definition Let (1.1) be an irregular equation with $k = 1$, Λ be the vector of eigenvalues of the corresponding matrix $A(0)$. Let (2.1) be its deformation depending continuously on a

nonnegative parameter ε , $f(t, 0) \equiv t^2$, $\alpha_0 + \alpha_1 \equiv 0$. The family (2.1) is said to be *generic*, if $\alpha_0(\varepsilon) \neq \alpha_1(\varepsilon)$ for $\varepsilon \neq 0$, and the line passing through $\alpha_0(\varepsilon)$ and $\alpha_1(\varepsilon)$ intersects each (k, Λ) -real dividing ray by angle bounded away from 0 uniformly in ε .

2.9 Definition Let (2.1) be a generic family (see the previous Definition), Λ be the corresponding eigenvalue tuple of $A(0) = A(0, 0)$. Let α_0, α_1 be the corresponding singular point families, V_i be the half-plane (depending on ε) containing α_i and bounded by the symmetry line of the segment $[\alpha_0, \alpha_1]$. The *sector associated to α_i* is a $(1, \Lambda)$ -good sector (see Definition 1.3) independent on ε that contains V_i for any ε small enough.

2.10 Remark In the previous Definition the sectors S_0, S_1 associated to α_0, α_1 respectively cover a punctured neighborhood of zero, so, the nonperturbed equation has a pair of Stokes operators (C_0, C_1) associated to this covering.

2.11 Theorem (see [6]). *Let (2.1) be a generic family of linear equations (see Definition 2.8), S_0, S_1, S'_0, S'_1 be respectively the corresponding associated sectors (see the previous Definition) and the domains (2.2). Then the statement of the previous Theorem remains valid. The same is true for Corollary 2.6. In more details, consider the eigenbases Z_ε^i on S'_i of the monodromy operators around the singular points $\alpha_i(\varepsilon)$ of the perturbed equation, $i = 0, 1$. Let $C_0(\varepsilon), C_1(\varepsilon)$ be the transition matrices (2.3) between them in the connected components of the intersection $S'_0 \cap S'_1$. Let C_0, C_1 be the Stokes matrices of the nonperturbed equation in the corresponding limit connected components of the intersection $S_0 \cap S_1$. Then for any $j = 0, 1$ and appropriately normalized monodromy eigenbases Z_ε^i (the normalization of Z_ε^0 (only) depends on the choice of j) $C_j(\varepsilon) \rightarrow C_j$, as $\varepsilon \rightarrow 0$.*

2.2 Stokes operators as limits of commutators of appropriate powers of the monodromy operators

The Stokes and monodromy operators act in different linear spaces: in the solution spaces of the nonperturbed (respectively, perturbed) equations. To formulate the statement from the title of the Subsection, let us firstly identify these solution spaces and specify the loops defining the monodromy operators.

Let (2.1) be a generic family of linear equations (in the sense of some of Definitions 2.1 and 2.8). Take a "base point" t_0 in the unit disc punctured at 0.

2.12 Remark The space of local solutions of a linear equation at a nonsingular point $t_0 \in \mathbb{C}$ is identified with the space of initial conditions at t_0 (which is common for the nonperturbed and the perturbed equations). This identifies the solution spaces of the latters. The space thus obtained will be denoted by H_{t_0} .

2.13 Remark Let (1.1) be an irregular equation with $k = 1$, Λ be the eigenvalue tuple of the corresponding matrix $A(0)$. Let S_0, S_1 be $(1, \Lambda)$ -good sectors covering a punctured neighborhood of zero in the t -line. Let C_0, C_1 be the Stokes operators (1.6) corresponding to the connected components of their intersection. Each operator C_i is well-defined in the space H_{t_0} of local solutions of (1.1) at any point t_0 lying in the corresponding component of the intersection $S_0 \cap S_1$.

Now let us define the monodromy operators acting in the previous space H_{t_0} .

2.14 Definition Let (2.1) be a generic family of linear equations (in the sense of one of Definitions 2.1 and 2.8), $\alpha_i(\varepsilon)$, $i = 0, 1$, be its singularity families. Fix a point t_0 (independent on ε) disjoint from the line passing through $\alpha_0(\varepsilon)$ and $\alpha_1(\varepsilon)$ for any ε . Let l_i be a small circle centered at $\alpha_i(\varepsilon)$ whose closed disc is disjoint from $-\alpha_i(\varepsilon)$, $a_i = [t_0, \alpha_i] \cap l_i$, the segment $[t_0, a_i]$ be oriented from t_0 to a_i . Consider the closed path $\psi_i = [t_0, a_i] \circ l_i \circ [t_0, a_i]^{-1}$, $i = 0, 1$, which starts and ends at t_0 (in the case, when $k = 1$, $n = 2$, $\lambda_1 - \lambda_2 \in \mathbb{R}$, we choose $t_0 \in \mathbb{R}$, see Fig.3). Define $M_i : H_{t_0} \rightarrow H_{t_0}$ to be the corresponding monodromy operator of the perturbed equation.




Figure 3

We show that commutators of appropriate noninteger powers of the operators M_i (see the following Definition) tend to the Stokes operators.

2.15 Definition Let $d \in \mathbb{R}$, $M : H \rightarrow H$ be a linear operator in a finite-dimensional linear space having distinct eigenvalues. The d -th power of M is the operator having the same eigenlines, as M , whose corresponding eigenvalues are some values of the d -th powers of those of M .

Let S_0, S_1 be sectors in $\mathbb{C}$ with vertex at 0 covering a punctured neighborhood of 0. Their left (right) intersection component is the component of their intersection crossed while going from S_0 to S_1 in the counterclockwise (respectively, clockwise) direction.

2.16 Theorem Let (2.1) be a generic family of linear equations (in the sense of one of Definitions 2.1 and 2.8), $\alpha_i(\varepsilon)$, $i = 0, 1$, be its singularity families. Let S_i , $i = 0, 1$, be the corresponding associated sectors (see Definitions 2.4 or 2.9 respectively) forming a covering of a punctured neighborhood of zero, C_0, C_1 be the Stokes operators (1.6) of the nonperturbed equation corresponding to the left (respectively, right) component of the intersection $S_0 \cap S_1$ (see the previous paragraph). Let t_0 be a fixed point of unit disc lying in the left component of the intersection $S_0 \cap S_1$, H_{t_0} be the corresponding local solution space (see Remark 2.12). (Then the operator C_0 (C_1) acts in the space H_{t_0} (respectively, H_{-t_0} , see Remark 2.13.)) Let

$M_i : H_{\pm t_0} \rightarrow H_{\pm t_0}$ be the corresponding monodromy operators from Definition 2.14. Then for any pair of numbers $d_0, d_1 > 0$ such that $d_0 + d_1 < 1$

$$\begin{aligned} M_1^{-d_1} M_0^{d_0} M_1^{d_1} M_0^{-d_0} &\rightarrow C_0 \text{ in the space } H_{t_0}, \\ M_0^{-d_0} M_1^{d_1} M_0^{d_0} M_1^{-d_1} &\rightarrow C_1 \text{ in the space } H_{-t_0}, \text{ as } \varepsilon \rightarrow 0. \end{aligned}$$

Theorem 2.16 is proved in Section 3.

2.3 The case of higher Poincaré rank

Theorem 2.5 on convergence of the monodromy eigenbases to canonical solution bases is stated and proved in [6] for arbitrary irregular equation (for arbitrary Poincaré rank and dimension). It holds for any generic family (1.2) defined to satisfy the following conditions: 1) $\sum \alpha_i \equiv 0$, $f'_\varepsilon(0, 0) \neq 0$ (then $\alpha_i(\varepsilon) = a_i \varepsilon^{\frac{1}{k+1}}(1 + o(1))$, where the points a_i form a regular polygon centered at 0); 2) no one of the previous points a_i lies in a real dividing ray (see Definition 2.7), in other terms, no radial ray of α_i tends to a real dividing ray. To each singularity family α we put into correspondence a (k, Λ) - good sector S (similarly to Definition 2.9) so that the canonical branches in $S'_\varepsilon = S \setminus \bigcup_{i=0}^k [0, \alpha_i(\varepsilon)]$ of the corresponding monodromy eigenfunctions converge to the canonical solutions of the nonperturbed equation on S . In the case of higher Poincaré rank for some pairs of neighbor singularities of the perturbed equation the corresponding sectors cannot not be chosen intersected; then the corresponding transition operator between the monodromy eigenbases tends to a product of Stokes operators. Each Stokes matrix is contained in some of the previous limit products, and its elements are expressed as polynomials in the elements of the corresponding limit product. On the other hand, in dimension two there are always two pairs of neighbor singularity families such that for each singularity pair the corresponding sectors may be chosen intersected. Then the transition operator between the corresponding appropriately normalized monodromy eigenbases tends to the Stokes operator of the nonperturbed equation corresponding to the intersection of the sectors.

2.17 Example Consider the case, when $k = n = 2$. Then the perturbed equation has three singularities, and the number of $(2, \Lambda)$ - good sectors covering a punctured neighborhood of zero is equal to 4. One can prove the following version of Theorem 2.16.

Consider a generic deformation (1.2) of an irregular equation (1.1) with $k = n = 2$. Let α_0, α_1 be a pair of singularity families numerated counterclockwise and corresponding to intersected sectors (denote the latters by S_0 and S_1 respectively). Let $t_0 \in \mathbb{C} \setminus 0$ be a fixed (base) point lying between the radial rays of $\alpha_0(\varepsilon), \alpha_1(\varepsilon)$ for all ε . Let M_0, M_1 be the corresponding monodromy operators (see Definition 2.14). Let C be the Stokes operator corresponding to the intersection $S_0 \cap S_1$. Then for appropriate $d_0, d_1 \in \mathbb{R} \setminus 0$ (depending on the family of equations)

$$M_1^{-d_1} M_0^{d_0} M_1^{d_1} M_0^{-d_0} \rightarrow C, \text{ as } \varepsilon \rightarrow 0.$$

More precisely, there exist $s_i \in \mathbb{N}$, $l_0, l_1 > 0$ (depending on the family of equations but not on ε) such that the previous statement holds whenever d_0, d_1 satisfy the following system of inequalities:

$$\begin{cases} (-1)^{s_i} d_i > 0, \quad i = 0, 1 \\ l_0 d_0 + l_1 d_1 < 1 \end{cases}.$$

2.18 Remark The previous coefficients l_i depend on how close the radial rays of α_i , $i = 0, 1$, approach the real dividing rays: if the minimal angle between the radial ray of α_i and some real dividing ray is small, then the corresponding coefficient l_i should be chosen large enough (hence, the corresponding exponent d_i should be taken small enough).

The author believes that the previous statements extend to the general case of arbitrary Poincaré rank and dimension.

3 Convergence of the commutators to Stokes operators. Proof of Theorem 2.16

Firstly we prove Theorem 2.16 in the case, when $k = 1$, $n = 2$. Its proof for the case of $k = 1$ and arbitrary n is analogous: the modifications needed will be discussed in Subsection 3.4

Thus, from now on we consider that $k = 1$, $n = 2$, until the contrary will be specified. Without loss of generality we assume that $\Lambda = (\lambda_1, \lambda_2) = (1, -1)$.

3.1 Properties of the monodromy and the transition operators. The plan of the proof of Theorem 2.16

Let us prove the convergence of the first commutator from Theorem 2.16; the proof of the convergence of the second commutator is analogous.

Thus, from now on we assume that the base point t_0 lies in the left component of the intersection $S_0 \cap S_1$, and one can put $t_0 = -\frac{1}{2}$.

3.1 Definition Consider a linear diagonalizable operator acting on $\mathbb{C}^2$ with eigenvalues of distinct modules. Its *projective multiplier* is the ratio of its eigenvalue with the lower module over that with the higher module. Its *projectivization* is the Möbius transformation $\overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ induced by its action and the tautological projection $\mathbb{C}^2 \setminus 0 \rightarrow \mathbb{P}^1 = \overline{\mathbb{C}}$.

3.2 Remark In the conditions of the previous Definition the projectivization is a hyperbolic transformation (see [1] and Definition 4.6 in Section 4); in particular, it has an attracting fixed point. The projective multiplier is well-defined and its module is always less than 1. It is equal to the multiplier of the projectivization at its attracting fixed point.

Let us write down the monodromy operators in the eigenbase of M_0 (which converges to the canonical solution base of the nonperturbed equation on S_0). Then the matrix of M_0 is diagonal: denote it

$$\Lambda_0(\varepsilon) = \text{diag}(\lambda_{01}, \lambda_{02})(\varepsilon).$$

By Corollary 2.6, the matrix of M_1 is

$$M_1 = C(\varepsilon)\Lambda_1(\varepsilon)C^{-1}(\varepsilon), \quad C(\varepsilon) \rightarrow C_0, \quad \text{as } \varepsilon \rightarrow 0, \quad \Lambda_1(\varepsilon) = \text{diag}(\lambda_{11}, \lambda_{12})(\varepsilon). \quad (3.1)$$

The transition matrix $C(\varepsilon)$ tends to the Stokes matrix C_0 , which is lower-triangular. Thus, the upper-triangular element of $C(\varepsilon)$ (denoted by $u(\varepsilon)$) tends to 0.

First of all we find the asymptotics of the eigenvalues λ_{ij} of M_i :

3.3 Proposition *Let (2.1) be a generic family of linear equations (see Definition 2.1), $t_0 = -\frac{1}{2}$, M_i be the monodromy operators of the perturbed equation from Definition 2.14, $f_{i1,\varepsilon}$, $f_{i2,\varepsilon}$ be their basic eigenfunctions, λ_{i1} , λ_{i2} be the corresponding eigenvalues. Then*

$$\lambda_{01}, \lambda_{12} \rightarrow \infty, \quad \lambda_{02}, \lambda_{11} \rightarrow 0,$$

$$\ln \lambda_{01} = -(1 + o(1)) \ln \lambda_{02} = -(1 + o(1)) \ln \lambda_{11} = (1 + o(1)) \ln \lambda_{12}, \quad \text{as } \varepsilon \rightarrow 0.$$

3.4 Corollary *In the conditions of the previous Proposition the projective multipliers of M_0 and M_1 are equal respectively to*

$$\mu_0 = \frac{\lambda_{02}}{\lambda_{01}}, \quad \mu_1 = \frac{\lambda_{11}}{\lambda_{12}}; \quad \mu_i \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0,$$

$$\ln \mu_0 = (1 + o(1)) \ln \mu_1, \quad \text{as } \varepsilon \rightarrow 0.$$

Proof of Proposition 3.3. It follows from definition that $\ln \lambda_{01} = (1 + o(1)) \frac{2\pi i}{\alpha_0 - \alpha_1}$. The real part of the right-hand side of the previous formula is positive and tends to infinity (since $\text{Im}(\alpha_0 - \alpha_1) > 0$ by assumption, and $\alpha_i \rightarrow 0$), which implies that $\lambda_{01} \rightarrow \infty$. The similar formulas written for all the λ_{ij} prove the rest of the statements of the Proposition. $\square$

Let $d_0, d_1 > 0$, $d_0 + d_1 < 1$,

$$\widetilde{M}_i = M_i^{d_i}, \quad \widetilde{\Lambda}_i = \Lambda_i^{d_i}, \quad i = 0, 1.$$

We prove that

$$\widetilde{M}_1^{-1} \widetilde{M}_0 \widetilde{M}_1 \widetilde{M}_0^{-1} \rightarrow C_0.$$

By definition and (3.1), the matrix of the previous commutator in the eigenbase of M_0 is

$$\widetilde{M}_1^{-1} \widetilde{M}_0 \widetilde{M}_1 \widetilde{M}_0^{-1} = C(\varepsilon) \widetilde{\Lambda}_1^{-1} C^{-1}(\varepsilon) \widetilde{\Lambda}_0 C(\varepsilon) \widetilde{\Lambda}_1 C^{-1}(\varepsilon) \widetilde{\Lambda}_0^{-1}. \quad (3.2)$$

Let $u(\varepsilon)$ be the upper-triangular element of the transition matrix $C(\varepsilon)$, $\mu_1(\varepsilon)$ be the projective multiplier of M_1 . For the proof of the convergence to $C_0 = \lim C(\varepsilon)$ of the previous commutator we firstly prove that

$$u = O(\mu_1), \quad \text{as } \varepsilon \rightarrow 0. \quad (3.3)$$

More precisely, we show in the next Subsection that $u = -(c_1 + o(1))\mu_1$, where c_1 is the upper-triangular element of the other Stokes matrix C_1 .

Let $\nu_0 = \mu_0^{d_0}$, $\nu_1 = \mu_1^{d_1}$ be the projective multipliers of the operators $\widetilde{M}_0$, $\widetilde{M}_1$ respectively. Formula (3.3) together with the previous Corollary and the condition $d_0 + d_1 < 1$ imply that

$$u(\varepsilon) = o(\nu_0 \nu_1), \quad \text{as } \varepsilon \rightarrow 0. \quad (3.4)$$

Using (3.4) we show (in the next Lemma proved in 3.3) that if we eliminate subsequently the terms $C^{\pm 1}(\varepsilon)$ in (3.2) (from the right to the left) except for the left $C(\varepsilon)$, then on each step the asymptotics of the modified expression (3.2) remains the same: the modified expression can be obtained from the initial one by composing it with an operator tending to the identity. At the last step the final modified expression will be just $C(\varepsilon)$, which tends to C_0 . This will prove the convergence of (3.2) to C_0 .

Now the convergence of the commutator (3.2) is implied by the following Lemma (modulo (3.3)).

3.5 Lemma *Let $\widetilde{M}_0, \widetilde{M}_1$ be two families of two-dimensional complex diagonalizable linear operators depending on a positive parameter ε . Let $\widetilde{\Lambda}_i = (\lambda_{i1}, \lambda_{i2})$ be the (diagonal) matrices of $\widetilde{M}_i$, $i = 0, 1$, in their eigenbases. Let $C(\varepsilon)$ be the transition matrix between the eigenbases, more precisely, in the eigenbase of $\widetilde{M}_0$ one have $\widetilde{M}_1 = C(\varepsilon)\widetilde{\Lambda}_1 C^{-1}(\varepsilon)$. Let the eigenbases converge to some bases in the space so that the corresponding transition matrix $C(\varepsilon)$ tends to a unipotent lower-triangular matrix (denoted by C_0). Let*

$$\nu_0 = \frac{\lambda_{02}}{\lambda_{01}} \rightarrow 0, \quad \nu_1 = \frac{\lambda_{11}}{\lambda_{12}} \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0. \quad (3.5)$$

Let ν_0, ν_1 and the upper-triangular element $u(\varepsilon)$ of the matrix $C(\varepsilon)$ satisfy (3.4). Then

$$\widetilde{M}_1^{-1} \widetilde{M}_0 \widetilde{M}_1 \widetilde{M}_0^{-1} \rightarrow C_0, \quad \text{as } \varepsilon \rightarrow 0.$$

The Lemma will be proved in Subsection 3.3.

Proof of Theorem 2.16 modulo (3.3) and Lemma 3.5. The operators $\widetilde{M}_i = M_i^{d_i}$ satisfy the conditions of the previous Lemma: all the conditions follow from the previous Proposition, Corollary and (3.3). This together with the Lemma implies the convergence of the first commutator in Theorem 2.16. This proves Theorem 2.16 modulo (3.3) and Lemma 3.5. $\square$

3.2 The upper-triangular element of the transition matrix. Proof of (3.3)

We prove the following more precise version of (3.3).

3.6 Lemma *Let (2.1) be a generic family of linear equations (see Definition 2.1), α_i be its singularity families, S_i be the corresponding sectors (see Definition 2.4) chosen to cover a punctured neighborhood of zero, S'_i be the corresponding domains from (2.2). Let C_0, C_1 be the Stokes matrices (1.6) of the nonperturbed equation (corresponding to the left (respectively, right) component of the intersection $S_0 \cap S_1$),*

$$C_0 = \begin{pmatrix} 1 & 0 \\ c_0 & 1 \end{pmatrix}, \quad C_1 = \begin{pmatrix} 1 & c_1 \\ 0 & 1 \end{pmatrix}, \quad \text{see Example 1.7.}$$

Let M_i be the monodromy operator of the perturbed equation around $\alpha_i(\varepsilon)$ acting in the space of solutions on S'_i . Let Z_ε^i be the (fundamental matrix of) its eigenbase. Let $C_0(\varepsilon)$ be the transition matrix (2.3) between the bases Z_ε^i in the left component of the intersection $S'_0 \cap S'_1$. Let the previous eigenbases be normalized to converge so that $C(\varepsilon) \rightarrow C_0$ (see Corollary 2.6):

$$C_0(\varepsilon) = \begin{pmatrix} 1 + o(1) & u(\varepsilon) \\ c_0 + o(1) & 1 + o(1) \end{pmatrix}, \quad u(\varepsilon) \rightarrow 0.$$

Let $\lambda_{11}, \lambda_{12}$ be the eigenvalues of M_1 , $\mu_1 = \frac{\lambda_{11}}{\lambda_{12}}$ be the corresponding projective multiplier. Then the upper-triangular element $u(\varepsilon)$ of the matrix $C_0(\varepsilon)$ has the asymptotics

$$u(\varepsilon) = (-c_1 + o(1))\mu_1^{-1}, \quad \text{as } \varepsilon \rightarrow 0, \quad (3.6)$$

where c_1 is the upper-triangular element of the Stokes matrix C_1 .

Proof The transition matrix $C_0(\varepsilon)$, $Z_\varepsilon^1 = Z_\varepsilon^0 C_0(\varepsilon)$, compares the monodromy eigenbases in the left component of the intersection $S'_0 \cap S'_1$, in particular, on a real interval in $\mathbb{R}_-$. It is not changed, when we extend the basic solutions analytically from $\mathbb{R}_-$ to $\mathbb{R}_+$ along the real line. Denote $Z_{\varepsilon,+}^i$ the corresponding branch on $\mathbb{R}_+$ of the extended fundamental matrix Z_ε^i , $i = 0, 1$. It follows from definition that $Z_{\varepsilon,+}^1$ is obtained from $Z_\varepsilon^1|_{\mathbb{R}_+}$ by applying the inverse monodromy operator M_1^{-1} :

$$Z_{\varepsilon,+}^1 = Z_\varepsilon^1|_{S'_1} M_1^{-1}; \text{ the matrix } M_1 \text{ is diagonal.} \quad (3.7)$$

On the other hand, we can choose a renormalization of the eigenbase $Z_{\varepsilon,+}^0$ by multiplication of the basic solutions by constants (i.e., changing it to $Z_{\varepsilon,+}^0 L(\varepsilon)$, $L(\varepsilon) = \text{diag}(l_1(\varepsilon), l_2(\varepsilon))$ is some family of diagonal matrices) so that in the right component of the intersection $S'_0 \cap S'_1$ the transition matrix $C_1(\varepsilon)$ between $Z_{\varepsilon,+}^0 L(\varepsilon)$ and Z_ε^1 tends to the Stokes matrix C_1 :

$$Z_{\varepsilon,+}^0 L(\varepsilon) = Z_\varepsilon^1|_{S'_1} C_1(\varepsilon), \quad C_1(\varepsilon) \rightarrow C_1.$$

By definition, $Z_{\varepsilon,+}^1 = Z_{\varepsilon,+}^0 C_0(\varepsilon)$. Substituting the latter and (3.7) to the previous formula yields

$$C_0(\varepsilon) = L(\varepsilon) C_1^{-1}(\varepsilon) M_1^{-1}. \quad (3.8)$$

The matrices $C_i(\varepsilon)$ tend to the Stokes matrices C_i , which are unipotent. The matrices $L(\varepsilon)$, M_1 are diagonal and depend on ε . This implies that

$$L(\varepsilon) = M_1(1 + o(1)), \text{ as } \varepsilon \rightarrow 0.$$

This together with (3.8) implies (3.6). $\square$

3.3 Commutators of operators with asymptotically common eigenline. Proof of Lemma 3.5

In the proof of Lemma 3.5 we use the following Proposition.

3.7 Proposition *Let $C(\varepsilon)$ be a family of two-dimensional matrices depending on a parameter $\varepsilon \geq 0$ and converging to a unipotent lower-triangular matrix, as $\varepsilon \rightarrow 0$. Let $u = u(\varepsilon)$ be the upper-triangular element of $C(\varepsilon)$ (thus, $u(\varepsilon) \rightarrow 0$). Let $\Lambda(\varepsilon) = \text{diag}(\lambda_1(\varepsilon), \lambda_2(\varepsilon))$ be a family of diagonal matrices depending on $\varepsilon > 0$ such that*

$$\nu = \frac{\lambda_1}{\lambda_2} \rightarrow 0, \quad u = o(\nu), \text{ as } \varepsilon \rightarrow 0. \quad (3.9)$$

Then

$$\Lambda^{-1}(\varepsilon) C(\varepsilon) \Lambda(\varepsilon) \rightarrow Id, \text{ as } \varepsilon \rightarrow 0. \quad (3.10)$$

Proof The diagonal elements of the matrix in (3.10) are equal to those of $C(\varepsilon)$, and thus, tend to 1. Its lower-triangular element tends to 0: it is equal to that of $C(\varepsilon)$ (which tends to a finite limit) times ν (which tends to 0 by (3.9)). Its upper-triangular element, which is equal to $u\nu^{-1}$, tends to 0 by (3.9). This proves (3.10). $\square$

Consider the commutator (3.2):

$$C(\varepsilon) \tilde{\Lambda}_1^{-1} C^{-1}(\varepsilon) \tilde{\Lambda}_0 C(\varepsilon) \tilde{\Lambda}_1 C^{-1}(\varepsilon) \tilde{\Lambda}_0^{-1}. \quad (3.11)$$

Using the previous Proposition, we firstly "kill" the right $C^{-1}(\varepsilon)$: we show that expression (3.11) is equal to

$$C(\varepsilon)\tilde{\Lambda}_1^{-1}C^{-1}(\varepsilon)\tilde{\Lambda}_0C(\varepsilon)\tilde{\Lambda}_1\tilde{\Lambda}_0^{-1}(Id + o(1)). \quad (3.12)$$

Then we kill similarly the right $C(\varepsilon)$ in (3.12) and the remaining $C^{-1}(\varepsilon)$. Finally we get that the initial commutator is equal to $C(\varepsilon)$ times the commutator of diagonal matrices (which is identity) times $(Id + o(1))$. This implies that (3.2) tends to $C_0 = \lim C(\varepsilon)$.

The first step: killing of C^{-1} . Let

$$Q(\varepsilon) = \tilde{\Lambda}_0 C^{-1} \tilde{\Lambda}_0^{-1}.$$

By definition, expression (3.11) is equal to

$$C(\varepsilon)\tilde{\Lambda}_1^{-1}C^{-1}(\varepsilon)\tilde{\Lambda}_0C(\varepsilon)\tilde{\Lambda}_1\tilde{\Lambda}_0^{-1}Q(\varepsilon).$$

It suffices to show that $Q(\varepsilon) \rightarrow Id$. This follows from the previous Proposition applied to the families of matrices $C^{-1}(\varepsilon)$ and $\Lambda(\varepsilon) = \tilde{\Lambda}_0^{-1}(\varepsilon)$: these families satisfy the conditions of the previous Proposition. Indeed, by (3.5), $\nu = \nu_0 \rightarrow 0$. The upper-triangular element of C^{-1} (denoted by $\tilde{u}$) is $\tilde{u} = O(u) = o(\nu_0\nu_1) = o(\nu_0)$ by (3.4). This proves (3.9) for $\tilde{u}$. The conditions of the Proposition are checked. Thus, by (3.10), $Q(\varepsilon) \rightarrow Id$.

The second step: killing of the right C in (3.12). It repeats the previous discussions with the families $C(\varepsilon)$ and $\Lambda(\varepsilon) = \tilde{\Lambda}_1\tilde{\Lambda}_0^{-1}$.

The third step: killing of the left C^{-1} . Done analogously by applying the previous Proposition to the matrix families C^{-1} and $\tilde{\Lambda}_1$. Lemma 3.5 is proved.

3.4 Convergence of the commutators to Stokes operators: the higher-dimensional case

The proof of Theorem 2.16 in higher dimensions repeats that in the two-dimensional case with some changes specified below.

Let (2.1) be a generic family of equations, α_i be its singularity families, $i = 0, 1$, S_i , be the corresponding associated sectors. Consider their "left intersection component" that is crossed while going counterclockwise from S_0 to S_1 . Let t_0 be a point lying in this component. Let H_{t_0} be the corresponding local solution space, $M_i : H_{t_0} \rightarrow H_{t_0}$ be the corresponding monodromy operators (see Definition 2.14). Let Z_ε^0 be the eigenbase of the monodromy operator M_0 , where the eigenfunctions are taken in the order of decreasing of the modules of the corresponding eigenvalues (it appears that these modules are really distinct, see the next Proposition). Let Z_ε^1 be that of M_1 , and the order of the eigenfunctions coincide with the order of increasing of the modules of the eigenvalues. Let $C(\varepsilon)$ be the transition matrix between them:

$$Z_\varepsilon^1 = Z_\varepsilon^0 C(\varepsilon).$$

Let C_0, C_1 be the Stokes matrices of the nonperturbed equation in the left (respectively, right) connected component of the intersection $S_0 \cap S_1$.

3.8 Proposition *In the above conditions for any ε small enough each monodromy operator M_i , $i = 0, 1$, has distinct eigenvalues (denote them $\lambda_{i1}, \dots, \lambda_{in}$). Moreover, for any $j, k = 1, \dots, n$, $j < k$, one has $\frac{\lambda_{0j}}{\lambda_{0k}} \rightarrow \infty$, $\frac{\lambda_{1j}}{\lambda_{1k}} \rightarrow 0$, as $\varepsilon \rightarrow 0$. Appropriately normalized eigenbases*

Z_ε^0 and Z_ε^1 converge to canonical solution bases of the nonperturbed equation in S_0 and S_1 respectively. Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of the matrix $A(0)$ numerated in the order of increasing of the values $\operatorname{Re} \frac{\lambda_j}{i\alpha_0(\varepsilon)}$. The numeration of each (converging) monodromy eigenbase corresponds to the numeration of the limit canonical solution base by the previous eigenvalues λ_j .

In dimension two the Proposition follows from Proposition 3.3. In higher dimension its proof is analogous to that of Proposition 3.3.

The Stokes matrices C_0 and C_1 are lower- (respectively, upper-) triangular. This is implied by the last statement of the previous Proposition and the following well-known fact.

3.9 Proposition (see, e.g., [9]). Let $k, n \in \mathbb{N}$, $n \geq 2$, (1.1) be an irregular equation, $\Lambda = (\lambda_1, \dots, \lambda_n)$ be the eigenvalues of the corresponding matrix $A(0)$. Let S_0, S_1 be a pair of intersected (k, Λ) -good sectors. Let there exist a $t \in S_0 \cap S_1$ such that the sequence of the values $\operatorname{Re} \frac{\lambda_j}{ik}$, $j = 1, \dots, n$, increases. Consider the canonical sectorial solution bases of (1.1) numerated by λ_j . Then the Stokes matrix of (1.1) in the connected component containing t of the intersection $S_0 \cap S_1$ is lower-triangular. In the case of the inverse order of the eigenvalues it is upper-triangular.

The point $t = i\alpha_0(\varepsilon)$ satisfies the conditions of the previous Proposition with $k = 1$ (the last statement of Proposition 3.8). Hence, by Proposition 3.9, the Stokes matrix C_0 is lower-triangular and C_1 is upper-triangular.

Let

$$C(\varepsilon) = (C_{ij}(\varepsilon)), \quad C'_1 = C_1^{-1} = (C'_{1,ij}).$$

Formula (3.6) of Lemma 3.6 extends to higher dimension as follows:

$$C_{jk}(\varepsilon) = (C'_{1,jk} + o(1)) \left(\frac{\lambda_{1j}}{\lambda_{1k}} \right), \quad \text{as } \varepsilon \rightarrow 0. \quad (3.13)$$

The proof of (3.13) repeats that of (3.6) in Subsection 3.2.

Let $d_0, d_1 > 0$, $d_0 + d_1 < 1$, $\tilde{\lambda}_{ij} = \lambda_{ij}^{d_i}$, $i = 0, 1$, $j = 1, \dots, n$. One has

$$C_{jk}(\varepsilon) = O\left(\frac{\tilde{\lambda}_{1j}}{\tilde{\lambda}_{1k}} \frac{\tilde{\lambda}_{0k}}{\tilde{\lambda}_{0j}}\right), \quad \text{as } \varepsilon \rightarrow 0. \quad (3.14)$$

Formula (3.14) follows from (3.13), the inequality $d_0 + d_1 < 1$ and the asymptotic formula $\ln \lambda_{0j} = -(1 + o(1)) \ln \lambda_{1j}$, $j = 1, \dots, n$, which is proved analogously to Proposition 3.3. As at the end of Subsection 3.1, Theorem 2.16 is implied by (3.14) and the following higher-dimensional analogue of Lemma 3.5.

3.10 Lemma Let $\tilde{M}_0, \tilde{M}_1$ be two families of n -dimensional diagonalizable linear operators depending on a positive parameter ε . Let $\tilde{\Lambda}_i = \operatorname{diag}(\lambda_{i1}, \dots, \lambda_{in})$ be the (diagonal) matrices of $\tilde{M}_i$ in their eigenbases. Let $C(\varepsilon) = (C_{jk}(\varepsilon))$ be the transition matrix between their eigenbases, more precisely, in the eigenbase of $\tilde{M}_0$ the matrix of $\tilde{M}_1$ is $C(\varepsilon)\tilde{\Lambda}_1 C^{-1}(\varepsilon)$. Let the eigenbases converge to some bases in the space so that the transition matrix $C(\varepsilon)$ converges to a unipotent lower-triangular matrix (denoted by C_0). Let for any $j < k$, $j, k = 1, \dots, n$,

$$\frac{\lambda_{0j}}{\lambda_{0k}} \rightarrow \infty, \quad \frac{\lambda_{1j}}{\lambda_{1k}} \rightarrow 0, \quad \text{as } \varepsilon \rightarrow 0.$$

Let in addition the asymptotic formula (3.14) hold. Then

$$\widetilde{M}_1^{-1} \widetilde{M}_0 \widetilde{M}_1 \widetilde{M}_0^{-1} \rightarrow C_0, \text{ as } \varepsilon \rightarrow 0.$$

The proof of the Lemma repeats that of Lemma 3.5 with obvious changes.

4 Generic divergence of monodromy operators along degenerating loops

In the present Section we consider only *two-dimensional irregular equations with Poincaré rank $k = 1$* and their generic deformations. As before, without loss of generality we assume that $\lambda_1 - \lambda_2 \in \mathbb{R}_+$, $\text{Im } \alpha_0 > 0$, $\text{Im } \alpha_1 < 0$.

Let (2.1) be a generic family of linear equations, α_i , $i = 0, 1$, be its singularity families, S_0 , S_1 be the corresponding associated sectors forming a covering of a punctured neighborhood of 0. Let $t_0 \in \mathbb{R}_-$ be arbitrary fixed base point independent on ε . Let $M_0 = M_0(\varepsilon)$, $M_1 = M_1(\varepsilon)$ be the corresponding monodromy operators of the perturbed equation (see Definition 2.14).

Consider the circle centered at 0 and passing through t_0 with the counterclockwise orientation (it bounds a disc containing both singularities of the perturbed equation for any small ε). The monodromy operator along the previous circle is called the *complete monodromy*.

4.1 Remark The complete monodromy of the perturbed equation in a generic family (2.1) converges to the monodromy of the nonperturbed equation along the counterclockwise circuit. In the previous conditions the complete monodromy is equal to $M_0 M_1$.

In the present Section we state and prove the Theorem saying that for any generic deformation (2.1) of a typical equation (1.1) (see the next Definition) each word (1.3) with integer exponents tends to infinity in GL_n , except for the powers $(M_0 M_1)^k$ of the complete monodromy.

4.1 The statement of the divergence Theorem

4.2 Definition Let (1.1) be an irregular equation, as at the beginning of the paper, $t_0 \in \mathbb{C} \setminus 0$ be arbitrary fixed base point, $M : H_{t_0} \rightarrow H_{t_0}$ be the counterclockwise monodromy operator around zero. Consider some branches at t_0 of all the sectorial canonical solutions of (1.1) as elements of H_{t_0} and take the collection of the complex lines in H_{t_0} generated by them. The equation is said to be *typical*, if for any $k \in \mathbb{Z} \setminus 0$ no line from the previous collection is transformed by M^k to another line from the same collection.

4.3 Remark The definition of typical equation does not depend on the choice of the base point and the branches of the canonical solutions. The condition that an equation (1.1) is typical is equivalent to a countable number of polynomial inequalities on the formal monodromy eigenvalues and the elements of the Stokes matrices.

4.4 Remark Both Stokes operators of a typical equation are nontrivial.

Each monodromy operator word (1.3) can be rewritten as

$$M_{j_n}^{s_n} \dots M_{j_1}^{s_1}, \quad s_i = \pm 1, \quad M_{j_k}^{s_k} M_{j_{k-1}}^{s_{k-1}} \neq 1 \text{ for any } k = 2, \dots, n, \quad (4.1)$$

where $n = \sum_{i=1}^l |d_i|$ in the notations of (1.3). We consider those words (4.1) that do not coincide literally with powers of the complete monodromy.

4.5 Definition A word (4.1) is said to be *reduced*, if it does not coincide (literally) with neither $M_0 M_1 \dots M_0 M_1$, nor $M_1^{-1} M_0^{-1} \dots M_1^{-1} M_0^{-1}$.

4.6 Theorem Let (1.1) be a typical equation (see Definition 4.2), (2.1) be its generic deformation, $t_0, M_0, M_1 : H_{t_0} \rightarrow H_{t_0}$, be as at the beginning of the Section. Then any monodromy operator given by a reduced word (4.1) tends to infinity (together with its projectivization, see Definition 3.1), as $\varepsilon \rightarrow 0$.

4.2 Projectivization. The scheme of the proof of Theorem 4.6

Instead of invertible linear operators $\mathbb{C}^2 \rightarrow \mathbb{C}^2$ we will consider their projectivizations, which are Möbius transformations $\overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$. Denote m_0, m_1 the projectivizations of the monodromy operators $M_0, M_1 : \mathbb{C}^2 \rightarrow \mathbb{C}^2$. We show that any reduced word $\tilde{m} = m_{j_n}^{s_n} \dots m_{j_1}^{s_1}$ tends to infinity in the Möbius group, as $\varepsilon \rightarrow 0$. This will prove the Theorem.

Recall the following

4.7 Definition (see [1]). A Möbius transformation $m : \overline{\mathbb{C}} \rightarrow \overline{\mathbb{C}}$ is said to be *hyperbolic*, if it has one repelling fixed point (then there is a unique attracting fixed point and each orbit except for the repeller tends to the attractor). A hyperbolic transformation with repeller a and attractor b will be presented as the picture at Fig.4




Figure 4

In the proof of the divergence of a reduced word $\tilde{m}$ of the projectivizations we use their following properties.

4.8 Proposition Let (2.1), S_i, t_0, M_i be as at the beginning of the Section (the nonperturbed equation is not necessarily typical), m_i be the projectivizations of M_i , $i = 0, 1$ (see Definition 3.1). Let $Z_\varepsilon^i = (f_{i1,\varepsilon}, f_{i2,\varepsilon})$ be the eigenbases of M_i , (f_{i1}, f_{i2}) be the sectorial canonical solution bases on S_i of the nonperturbed equation, $i = 0, 1$. Let $p_{ij,\varepsilon}, p_{ij} \in \overline{\mathbb{C}}$ be the tautological projection images of $f_{ij,\varepsilon}$ and f_{ij} respectively. Then m_i are hyperbolic transformations (see the previous Definition) with fixed points $p_{ij,\varepsilon}$: $p_{02,\varepsilon}, p_{01,\varepsilon}$ are respectively the repelling and

attracting fixed points of m_0 ; $p_{11,\varepsilon}$, $p_{12,\varepsilon}$ are respectively the repelling and attracting fixed points of m_1 .

Proof The Proposition follows from Proposition 3.3 □

4.9 Proposition *Let (2.1), S_i , t_0 , M_i be as at the beginning of the Section, p_{ij} , $p_{ij,\varepsilon}$ be the tautological projection images of the canonical basic solutions of the nonperturbed equation and the eigenfunctions of M_i respectively (see the previous Proposition). Then*

$$p_{02} = p_{12}, p_{ij} = \lim_{\varepsilon \rightarrow 0} p_{ij,\varepsilon}, \text{ see Fig.5a,b.} \quad (4.2)$$




Figure 5

Proof The statements on the limits in (4.2) follow from Theorem 2.5. The coincidence of p_{02} and p_{12} in (4.2) follows from the lower-triangularity of the Stokes matrix C_0 (see Example 1.7). □

As it is shown below, Theorem 4.6 is implied by the two previous Propositions and the following Lemma.

4.10 Lemma *Let $p_{02} = p_{12}$, p_{01} , p_{11} be a given triple of distinct points in $\overline{\mathbb{C}}$. Let $m_i = m_i(\varepsilon)$, $i = 0, 1$, be two families of hyperbolic Möbius transformations depending on a positive parameter ε , μ_0 , μ_1 be the multipliers of their attractors. Let $p_{01,\varepsilon}$, $p_{12,\varepsilon}$ be respectively the attractors of m_0 and m_1 , $p_{02,\varepsilon}$, $p_{11,\varepsilon}$ be their repellers. Let $p_{ij,\varepsilon} \rightarrow p_{ij}$, $\mu_i \rightarrow 0$, as $\varepsilon \rightarrow 0$. Let in addition the product $m_0 m_1$ converge to a Möbius transformation m such that for any $k \in \mathbb{Z} \setminus 0$, $i = 0, 1$, $j = 1, 2$ the image $m^k p_{ij}$ coincide with no other p_{ls} . Then any reduced word $\tilde{m} = m_{j_n}^{s_n} \dots m_{j_1}^{s_1}$ (see Definition 4.5) tends to infinity.*

The projectivizations of the monodromy operators satisfy the conditions of the Lemma. Indeed, the multipliers μ_i tend to 0 by Corollary 3.4. The convergence $p_{ij,\varepsilon} \rightarrow p_{ij}$ follows from the previous Proposition. The product $m_0 m_1$ tends to the projectivization (denoted by m) of the monodromy of the nonperturbed equation. The inequalities $m^k p_{ij} \neq p_{ls}$ follow from typicality (see Definition 4.2). This together with the Lemma proves divergence of $\tilde{m}$. Theorem 4.6 is proved modulo the Lemma.

4.3 Divergence of word of projectivizations. Proof of Lemma 4.10

As it is shown below, Lemma 4.10 is implied by the following statement.

4.11 Lemma *In the conditions of the previous Lemma let $\tilde{m} = m_{j_n}^{s_n} \dots m_{j_1}^{s_1}$ be a reduced word such that*

$$m_{j_n}^{s_n} m_{j_{n-1}}^{s_{n-1}} \neq (m_0 m_1)^{\pm 1}. \quad (4.3)$$

Let $x \in \overline{\mathbb{C}}$ be arbitrary point such that

$$m^s x \neq p_{ij} \text{ for any } s = -n, \dots, n, \ i = 0, 1, \ j = 1, 2; \ m = \lim_{\varepsilon \rightarrow 0} m_0 m_1. \quad (4.4)$$

Then the image $\tilde{m}x$ converges to the limit of the attractor of $m_{j_n}^{s_n}$.

Let us prove Lemma 4.10 modulo Lemma 4.11. If the reduced word $\tilde{m}$ under consideration satisfies (4.3), then it tends to infinity. Indeed, by Lemma 4.11, the image $\tilde{m}x$ of a generic x tends to the attractor of $m_{j_n}^{s_n}$, hence, $\tilde{m} \rightarrow \infty$. Otherwise, $\tilde{m} = (m_0 m_1)^k m'$, where $k \in \mathbb{Z} \setminus 0$, m' is a word satisfying (4.3) of a length less than that of $\tilde{m}$. The new word m' tends to infinity by the previous statement. The product $m_0 m_1$ in the previous expression for $\tilde{m}$ has a finite limit. Hence, $\tilde{m}$ tends to infinity as well. Lemma 4.10 is proved.

Proof of Lemma 4.11. In the proof of Lemma 4.11 we use the following obvious

4.12 Proposition *Let $m' = m'(\varepsilon)$ be a family of hyperbolic Möbius transformations depending on a positive parameter ε , $m' \rightarrow \infty$, as $\varepsilon \rightarrow 0$, so that the attractor and the repeller of m' tend to distinct limits (hence, the multiplier of the attractor tends to 0). Then for any point $x \in \overline{\mathbb{C}}$ distinct from the limit of the repeller its image $m'x$ tends to the same limit, as the attractor. The same statement holds for arbitrary family $x(\varepsilon)$ of points bounded away from the limit of the repeller.*

We prove Lemma 4.11 by induction in the length n of the word. For $n = 1$ its statement is obvious. Suppose we have proved the Lemma for the words of any length less than a given n . Let us prove it for a word $\tilde{m} = m_{j_n}^{s_n} \dots m_{j_1}^{s_1}$ of the length n .

Without loss of generality we assume that $m_{j_n}^{s_n} = m_0$: the contrary case is treated analogously. Then by (4.3) and the inequality $m_{j_n}^{s_n} m_{j_{n-1}}^{s_{n-1}} \neq 1$ (see (4.1)),

$$m_{j_{n-1}}^{s_{n-1}} \neq m_1, m_0^{-1}, \text{ thus, } m_{j_{n-1}}^{s_{n-1}} = m_0 \text{ or } m_1^{-1}. \quad (4.5)$$

Consider the word

$$m' = m_{j_{n-1}}^{s_{n-1}} \dots m_{j_1}^{s_1} = m_{j_n}^{-s_n} \tilde{m}.$$

Suppose firstly that it satisfies (4.3). Then for any x satisfying (4.4) $m'x$ tends to the limit (denoted p_{ij}) of the attractor of $m_{j_{n-1}}^{s_{n-1}}$ by the induction hypothesis. The latter attractor can be either $p_{01,\varepsilon}$, or $p_{11,\varepsilon}$, which are the attractor of m_0 and the repeller of m_1 respectively. This follows from (4.5). Hence, the limit p_{ij} is either p_{01} , or p_{11} ; in both cases it does not coincide with the limit p_{02} of the repeller of m_0 . Therefore, the image $m_0 p_{ij}$ (and hence, $m_0(m'x) = \tilde{m}x$) tends to the same limit, as the attractor of m_0 (by the previous Proposition).

Now suppose that the word m' does not satisfy (4.3). Then

$$m' = (m_0 m_1)^k m'',$$

where $k \in \mathbb{Z} \setminus 0$, and m'' is a word satisfying (4.3) of a length less than that of m' . The induction hypothesis applied to m'' implies that for any x satisfying (4.4) its image $m''x$ tends to the limit of the attractor of the last left element of the word m'' , thus, to some p_{ij} . Therefore, $m'x \rightarrow m^k p_{ij}$. The image $m^k p_{ij}$ coincides with no p_{sl} (in particular, with the limit p_{02} of the repeller of m_0). This follows from the last condition of Lemma 4.10. Therefore, $m_0 m^k p_{ij}$ tends to the limit of the attractor of m_0 (and so does $m_0 m'x = \tilde{m}x$ by the previous Proposition). The induction step is over. Lemma 4.11 is proved. The proof of Lemma 4.10 is completed. $\square$

I am grateful to V.I. Arnold, Yu.S. Ilyashenko and A.A. Bolibrukh for stating the problems. I wish to thank them and V. Kleptsyn for their interest to the work and helpful remarks. The research was supported by part by CRDF grant RM1-2358-MO-02 and by Russian Foundation for Basic Research (RFFI) grant 02-02-00482.

References

- [1] Apanasov, B.N., Gusevskii, N.A., Krushkal, S.L., *Kleinian groups and uniformization in examples and problems*. Translated from the Russian by H. H. McFaden. Translation edited and with a preface by Bernard Maskit. Translations of Mathematical Monographs, 62. American Mathematical Society, Providence, RI, 1986.
- [2] Arnold, V.I., Ilyashenko, Yu. S., Ordinary differential equations. - In the volume *Dynamical Systems-1* of the series "Itogi nauki i tekhniki. Sovremennyye problemy matematiki. Fundamentalnyye napravleniya", VINITI publisher, Moscow (1985). - English translation in *Encyclopaedia Math. Sci.*, 1, *Dynamical systems*, I, 1-148, Springer, Berlin, 1988.
- [3] Balser, W.; Jurkat, W.B.; Lutz, D.A. Birkhoff invariants and Stokes' multipliers for meromorphic linear differential equations, *J. Math. Anal. Appl.* 71 (1979), No.1, 48-94.
- [4] Duval, A. Biconfluence et groupe de Galois, *J. Fac. Sci. Tokyo*, Sect. IA, Math. 38 (1991), No.2, p.211-223.
- [5] Garnier, R., Sur les singularités irrégulières des équations différentielles linéaires, *J. Math. Pures et Appl.*, 8^e série, 2 (1919), pp. 99-198.
- [6] Glutsyuk, A. Stokes operators via limit monodromy of generic perturbation, *J. Dyn. & Contr. Syst.*, vol. 5 (1999), No 1, 101-135.
- [7] Glutsyuk, A. Confluence of singular points and the nonlinear Stokes phenomena (in Russian), *Trudy Moskovskogo Matematicheskogo Obshchestva*, vol.62 (2001), pp 54-104. English translation in *Transactions of Moscow Math. Soc.*, 2001, 49-95.
- [8] Glutsyuk, A. Resonant confluence of singular points and Stokes phenomena, *to appear*.
- [9] Ilyashenko, Yu. S.; Khovanskii, A. G., Galois groups, Stokes operators and a theorem of Ramis. (Russian) *Funktsional. Anal. i Prilozhen.* 24 (1990), no. 4, 31-42, 96; translation in *Funct. Anal. Appl.* 24 (1990), no. 4, 286-296 (1991).

- [10] Jurkat, W.B.; Lutz, D.A.; Peyerimhoff, A. Birkhoff invariants and effective calculations for meromorphic linear differential equations, *J. Math. Anal. Appl.* 53 (1976), No.2, 438–470.
- [11] Jurkat, W.B.; Lutz, D.A.; Peyerimhoff, A. Birkhoff invariants and effective calculations for meromorphic linear differential equations, *Houston J. Math.* 2 (1976), no. 2, 207–238.
- [12] Mardesic, P., Roussarie, R., Rousseau, C., Modulus of analytic classification for unfoldings of generic parabolic diffeomorphisms, Prépublication No 292 (2002) du Laboratoire de Topologie et du Laboratoire Gevrey de Mathématique Physique, Université de Bourgogne., Dijon, France.
- [13] J.Martinet. Remarques sur la bifurcation nœud-col dans le domaine complexe, *Astérisque* No 150-151, 1987. P.131-149.
- [14] Ramis, J.-P., Phénomène de Stokes et filtration Gevrey sur le groupe de Picard-Vessiot, *C. R. Acad. Sci. Paris Sér. I Math.* 301 (1985), no. 5, 165–167.
- [15] Ramis, J.-P. Confluence et resurgence, *J. Fac. Sci. Tokyo, Sect. IA, Math.* 36 (1989), No.3, p.703-716.
- [16] Sibuya, Y., Stokes phenomena, *Bull. Amer. Math. Soc.*, **83** (1977), pp. 1075–1077.
- [17] Zhang, C., Quelques études en théorie des équations fonctionnelles et en analyse combinatoire, - Thèse, Institut de Recherche Mathématique Avancée, Univ. Louis Pasteur et CNRS (URA 01), 1994.

*CNRS, Unité de Mathématiques Pures et Appliquées, M.R.,
 École Normale Supérieure de Lyon
 46 allée d'Italie, 69364 Lyon Cedex 07 France*